













TOBACCO FACTS

Tobacco use	Types of tobacco use		Effects of tobacco on oral health
<p>  800 million men smoke.  200 million women smoke. </p> <hr/> <p> 600,000 individuals die each year from secondhand smoke: 156,000 men, 281,000 women and 166,000 children. </p> <hr/> <p>At least 300 million people use smokeless tobacco and 90% of these are in Southeast Asia.</p> <hr/> <p>In 2011, manufacturers spent about US\$9.5 billion on advertising cigarettes and smokeless tobacco.</p> <hr/> <p>Governments spend less than US\$1 billion on tobacco control each year.</p>	<p>Smoking</p> <p>cigarettes </p> <p>bidis </p> <p>kreteks </p> <p>pipes </p> <p>cigars </p> <p>waterpipes </p>	<p>Smokeless</p> <p>snuff, dry and moist </p> <p>chewing tobacco </p> <p>snus </p> <p>dissolvables </p>	<p>Increases risk of:</p> <ul style="list-style-type: none"> • oral cancer • smoker's palate • periodontal disease • premature tooth loss • gingivitis • staining • halitosis (bad breath) • loss of taste and smell

From *The Challenge of Oral Disease – A call for global action* by FDI World Dental Federation.
 Maps and graphics © Myriad Editions 2015